

# Wheat Seeding Rate Evaluation

Andy Kleinschmidt, AGNR Extension Educator- Van Wert County  
Gary Prill, Program Manager- Farm Focus Research- Van Wert County

## Objective

The objective of this study was to evaluate the yield response of one wheat variety to four different seeding populations ranging from 1.2 to 2.1 million seeds per acre.

## Background

---

Crop Year:	2008	Herbicide:	
Cooperator:	Farm Focus/Marsh Foundation	POST	Express 75DFat 0.33 oz/A +
County/Town:	Van Wert/Van Wert	(April 23)	MCPA Amine at 0.75 pt/A +
Soil Type:	Hoytville clay; Hoytville silty clay loam	Variety:	AgriPro Cooker- Branson
Drainage:	Non-systematic tile	Insecticide:	Mustang Max @ 3 oz/A on June 17 for armyworm
Previous Crop:	Soybeans	Row Width:	7.5 inches
Tillage:	No-till	Planting Rate:	Variable-see Methods
Soil Test(2005):	pH 6.7, P 47 ppm, K 156 ppm	Planting Date:	October 8, 2007
Fertilizer:	245 lb/A 13-9-18 surface broadcast (October 5)	Harvest Date:	July 7, 2008
	220 lb/A 37-0-0-8.5S broadcast topdress (April 17)		

---

## Methods

This study was set up with four different planting populations replicated four times in a randomized complete block design. The populations used in this trial were:

- 1) 1.2 million seeds/acre
- 2) 1.5 million seeds/acre
- 3) 1.8 million seeds/acre
- 4) 2.1 million seeds/acre

The study was planted on October 8, 2007 using a John Deere 750 no-till drill. The drill was calibrated for the proper seeding rate at each population based on the 11,500 seeds per pound seed count reported on the AgriPro Cooker Branson variety seed tag. Plot size was 27.5 feet wide by 1070 feet long. Plots with wheel tracks from the late season spraying of insecticide for armyworm control had 1 foot of width subtracted if there was one wheel track within a plot, and 2 feet subtracted if there were two wheel tracks within a plot. Not all plots had wheel tracks so this adjustment to plot width was required to offset wheat that was flattened down by this application.

Harvest populations were estimated by counting the number of heads in one foot of row at six separate locations within each plot. These counts were converted to heads per acre. Harvesting was accomplished with a John Deere 6620 combine equipped with a calibrated AgLeader PF3000 yield monitor. Plot weights were determined with a calibrated weigh wagon. Moistures were taken from the yield monitor average reading for each plot. All yields were adjusted to 13.5% moisture.

## Results

**Table 1.** Harvest moisture, yield, and head count means.

Treatment	Moisture	Yield	Head Count
	(%)	(bu/A)	(heads/A)
1.2 million seeds/acre	13.5	92.5	3,917,500
1.5 million seeds/acre	13.3	95.6	3,720,300
1.8 million seeds/acre	13.4	95.5	4,057,000
2.1 million seeds/acre	13.4	97.3	4,068,500
LSD (P=0.05)	NS	NS	NS
F-test	2.6	2.7	3.2
CV(%)	< 1	2.5	4.6

NS= not significant

## Summary

The results of this single year wheat seeding rate study suggest there were no statistical differences between the treatments for moisture, yield, or head count. The head counts taken in this study would suggest that lower seeding rates still were able to produce about the same number of heads per acre, and thus similar yields.

From an economic standpoint, each 300,000 increase in the seeding rate between the treatments increased seed cost an additional \$8.32 per acre. This is based on a 2007 seed cost of \$15.95 per 50 pound bag of seed. This would require an additional 1.4 bushel yield increase for each additional 300,000 seeds dropped based on a harvest time market price of \$6 per bushel.

According to Ohio State University agronomists, the optimum seeding rate is 1.2 to 1.6 million seeds per acre (18 to 24 seeds per foot of 7.5-inch row) when planting during the two weeks following the fly-safe date. During the third and fourth week after the fly-safe date, plant 1.6 to 2.0 million seeds per acre (24 to 30 seeds per foot of row). Fly-safe date for Van Wert County is September 26. Planting date for this trial was 12 days after the fly-free date.

A similar study conducted at Farm Focus in 2007 ([http://farmfocus.osu.edu/wheat\\_seeding\\_rate-07.pdf](http://farmfocus.osu.edu/wheat_seeding_rate-07.pdf)) indicated there was a significant difference in yield between 1.2 million seeds/acre and 2.0 million seeds/acre seeding rates. The 2007 study planting date was later at 17 days after fly-free date, and weather conditions in the fall of 2007 were much less conducive to stand establishment and tillering than those experienced in the fall of 2008.

## Acknowledgement

OSU Extension- Van Wert and Farm Focus express appreciation to AgriPro Cooker for supplying the seed for this study.

For additional information contact:

Gary Prill

1055 South Washington Street

Van Wert, OH 45891

419-238-1214

prill.1@osu.edu



Ohio State University Extension embraces human diversity and is committed to ensuring that all research and related educational programs are available to clientele on a nondiscriminatory basis without regard to race, color, religion, sex, age, national origin, sexual orientation, gender identity or expression, disability, or veteran status. This statement is in accordance with United States Civil Rights Laws and the USDA.

Keith L. Smith, Ph.D., Associate Vice President for Agricultural Administration and Director, Ohio State University Extension

TDD No. 800-589-8292 (Ohio only) or 614-292-1868